APPOINTMENT

CONVERSATION COMPANION

Getting the most out of your doctor appointments



Once you've started your prescribed treatment with RYSTIGGO®, you'll start to have more focused conversations with your healthcare provider. You'll want to dedicate part of your appointment time to talk about topics such as how you are feeling and how you are managing your generalized myasthenia gravis (gMG) with RYSTIGGO.





Track, Discuss, Plan

It's important to keep conversations on track during your appointments. The time with your healthcare provider may be brief, so you'll want to plan ahead and show up ready. You can use

this document to help you with tracking your symptoms, planning for future treatment with your healthcare provider, and preparing for what's next. After all, this is your time to learn more about your treatment and get your questions answered.



പ്പ് Track Your Symptoms

An important aspect of treatment with RYSTIGGO is symptom tracking. These questions can help you make sure you're on the same page with your healthcare provider about this essential task.

	Why is i	it important	for me to	track my	symptoms?
--	----------	--------------	-----------	----------	-----------

- How (and how often) do you recommend I track my symptoms?
- Should I track anything other than symptoms?
- How will we discuss my symptom tracking at my next appointment? What information will you be looking for?



For tracking tools and other resources, visit ucbONWARD.com/RYSTIGGO.

INDICATION

RYSTIGGO (rozanolixizumab-noli) is indicated for the treatment of generalized myasthenia gravis (gMG) in adult patients who are anti-acetylcholine receptor (AChR) or antimuscle-specific tyrosine kinase (MuSK) antibody positive.

IMPORTANT SAFETY INFORMATION

RYSTIGGO may increase the risk of infection and could cause aseptic meningitis.

Tell your healthcare provider right away if you have signs or

symptoms of an infection or meningitis during treatment with RYSTIGGO. Swelling and rash have also occurred. Tell your healthcare provider immediately about any undesirable reactions you experience after administration. The most common side effects of RYSTIGGO include headache, infections, diarrhea, fever, hypersensitivity reactions, and nausea.

Please see the Important Safety Information on page 3 and the accompanying full Prescribing Information.



Discuss How to Get the Most Out of RYSTIGGO

Whether you're just getting started or already making progress in your treatment plan, these questions can help you learn more about your prescribed treatment with RYSTIGGO.

 What else should I know about RYSTIGGO that I haven't asked? Why do I need to attend all of my infusion appointments? What side effects might I experience on RYSTIGGO, and what should I do if I experience them? What additional resources for gMG or RYSTIGGO do you recommend? Are at-home infusions possible for me? If so, how can I get my infusions done at home? 				
Plan Future Treatment Cycles With Your Healthcare Team				
These questions can help you and your doctor decide when to begin your next RYSTIGGO treatment cycle, if necessary.*				
Who should I call for questions or support between appointments?How often should we meet to discuss my RYSTIGGO treatment cycles?				
How will we decide if and when I need additional RYSTIGGO treatment cycles?				
What information do you need from me to make that decision?				

*Most people may require repeated cycles of RYSTIGGO to manage their gMG symptoms. Everyone responds to treatment differently, so it's important to monitor your progress. After your treatment cycle, work with your healthcare provider to determine if or when another cycle is needed.





INDICATION

RYSTIGGO is a prescription medicine used to treat adults with a disease called generalized myasthenia gravis (gMG) who are acetylcholine receptor (anti-AChR) antibody positive or muscle-specific tyrosine kinase (anti-MuSK) antibody positive.

IMPORTANT SAFETY INFORMATION

Infection: RYSTIGGO may increase the risk of infection. In clinical studies, the most common infections were upper respiratory tract infections, COVID-19, urinary tract infections, and herpes simplex infections. Your healthcare provider should check you for infections before starting and during treatment with RYSTIGGO. Tell your healthcare provider if you have any history of infections. Tell your healthcare provider right away if you have signs or symptoms of an infection during treatment with RYSTIGGO. Some of the signs and symptoms may include fever, chills, frequent and/or painful urination, cough, runny nose, wheezing, shortness of breath, fatigue, sore throat, excess phlegm, nasal discharge, back pain, and/or chest pain.

Aseptic Meningitis: RYSTIGGO could cause aseptic meningitis. Tell your healthcare provider right away if you develop any signs or symptoms of meningitis during treatment with RYSTIGGO such as severe headache, neck stiffness, drowsiness, fever, sensitivity to light, painful eye movements, nausea and vomiting.

Hypersensitivity Reactions: RYSTIGGO can cause swelling and rash. Your healthcare provider should monitor you during and after treatment and discontinue RYSTIGGO if needed. Tell your healthcare provider immediately about any undesirable reactions you experience after administration.

Before taking RYSTIGGO, tell your healthcare provider about all of your medical conditions, including if you have a history of infection or think you have an active infection or have received or are scheduled to receive a vaccine (immunization). The use of vaccines during RYSTIGGO treatment has not been studied, and the safety with live or live-attenuated vaccines is unknown. Administration of live or live-attenuated vaccines is not recommended during treatment with RYSTIGGO. Completion of age-appropriate vaccines according to vaccination guidelines before starting a new treatment cycle with RYSTIGGO is recommended.

Before taking RYSTIGGO, tell your healthcare provider if you are pregnant or plan to become pregnant, or are breastfeeding or plan to breastfeed.

Tell your healthcare provider about all the medicines you take, including prescription and over-the-counter medicines, vitamins, and herbal supplements.

The most common side effects of RYSTIGGO include headache, infections, diarrhea, fever, hypersensitivity reactions, and nausea.



These are not all of the possible side effects of RYSTIGGO. For more information, ask your healthcare provider or pharmacist. Tell your healthcare provider about any side effect that bothers you or that does not go away. Call your healthcare provider for medical advice about side effects. You are encouraged to report negative side effects of prescription drugs to the FDA. Visit www.fda.gov/medwatch or call 1-800-FDA-1088. You may also report side effects to UCB, Inc. by calling 1-844-599-CARE [2273].

Please see the full Prescribing Information and talk to your healthcare provider about your condition or your treatment. For more information, go to www.RYSTIGGO.com or call 1-844-599-2273.